

median age was 31 years (26-42 years) and 33 were women (52.4%).The median symptom duration of HAE was 14 years (6-23 years). The median onset age of HAE was 17 years (12–21 years). A total of 10 patients were being followed up under long-term prophylaxis, seven patients were on danazol, and three patients were on tranexamic acid.

Table I: Demographic characteristics of HAE patients.

Clinical Characteristics	Total (N = 63)
Female ,n (%)	33 (52.4)
Age (years) (median, IQR)	31 (26-42)
Body Mass Index (mean±SD)	26.38±4.08
Smoker,n (%)	13 (20.6)
Alcohol consumption,n (%)	7 (11.1)
Marriage status,n (%)	
Single	15 (23.8)
Married	45 (71.4)
Widow	1 (1.6)
Divorced	2 (3.2)
Education status,n (%)	
Illiterate	3 (4.8)
Literate	8 (12.7)
Primary	27 (42.9)
High school	17 (27)
University	8 (12.7)
Employment status,n (%)	
Employed	29 (46)
Unemployed	31 (49.2)
Retired	1 (1.6)
Student	2 (3.2)
HAE symptoms, n (%)	
Pre-earthquake	
Abdominal	51 (81)
Larenx	21 (33.3)
Face	26 (41.3)
Extremity	52 (82.5)
Genital	14 (22.2)
Post-earthquake	
Abdominal	40 (63.5)
Larenx	20 (31.7)
Face	19 (30.2)
Extremity	41 (65.1)
Genital	12 (19)
HAE attack number (mean±SD)	
Pre-earthquake	5.2 ± 2.6
Post-earthquake	9.2 ± 5.2
Long term prophylaxis,n (%)	
Danazol	7 (11.1)
Tranexamic acid	3 (4.8)
C1 esterase inhibitor	0 (0)
None	53 (84.1)
HAE type,n (%)	
Type 1	62 (98.4)
Type 2	1 (1.6)
C4 level (median, IQR) (mg/dl) (n:62)	4 (2-7)
C1 esterase inhibitor level (median, IQR) (mg/dl) (n:62)	5.77 (4-8.82)
C1 esterase inhibitory function (%) (n:1)	<10

HAE, hereditary angioedema; IQR, interquartile range; SD, standard deviations

Physical condition after the earthquake

The effect of an earthquake on the course of hereditary angioedema according to individual answers was worsened in 45 (71.4%) patients, and not affected in 18 (28.6%) patients. The median (IQR) attacks of HAE were 8 (7-10) after the earthquake. The most common symptoms after the earthquake included extremity edema (65.1%), abdominal pain (63.5%), larynx edema (31.7%), face edema (30.2.5), and genital edema (19%).

The use of icatibant after the earthquake was 61 (96.8%). The numbers of post-earthquake attacks and icatibant injections were significantly higher than the ones in the pre-earthquake period (p<0.001) (Figure 1). The use of C1 esterase inhibitors after the earthquake was 12 (19%). C1 esterase inhibitor use did not significantly change (p>0.05). The median (IQR) VAS10 for attack severity was 8 (6-9). The attacks in the post-earthquake period were more severe than those in the pre-earthquake period (p<0.001) (Figure 1).




Figure 1. (A) Median number of HAE attacks in pre- and post-earthquake, (B) The median VAS severity scores of HAE attacks in pre- and post-earthquake, (C) Median number of icatibant injections in pre- and post-earthquake, (D) Median number of c1 esteraseinhibitor injections in pre- and post-earthquake. Abbreviations: IQR, interquartile range; VAS, visual analogue scale; HAE, hereditary angioedema

Psychological condition after the earthquake

As shown in Table 2, DASS-Stress, DASS-Anxiety, DASS-Depression subscale analysis revealed that 25(39.7%), 37(58.7%) and 32(50.8%) patients

had abnormal stress, anxiety and depression status, respectively. DASS-Stress, DASS-Depression, DASS-21 total scores and FES scores were significantly higher in women than men. Although the rate of patients with worsened HAE during the earthquake was higher in women (60%) than men (40%), it was not statistically significant (p=0.06). DASS-Stress, DASS-Anxiety, DASS-Depression and FES scores were significantly higher in patients with worsened HAE during earthquake than others (Table 3). Positive correlations between the number, and the severity of HAE attacks and DASS-21 anxiety, stress and depression were observed in Table 4.

Table II: The frequencies of DASS-21 and FES scale analysis in HAE patients after earthquake

	N =63	Female (n=33)	Male (n=30)	P value
DASS-Stress analysis, n (%)				
Normal	38 (60.3)	16 (48.5)	22 (73.3)	0.06
Mild	6 (9.5)	5 (15.2)	1 (3.3)	
Moderate	6 (9.5)	2 (6.1)	4 (13.3)	
Severe	10 (15.9)	9 (27.3)	1 (3.3)	
Very severe	3 (4.8)	1 (3)	2 (6.7)	
DASS-Anxiety analysis, n (%)				
Normal	26 (41.3)	11 (33.3)	15 (50)	0.12
Mild	8 (12.7)	5 (15.2)	3 (10)	
Moderate	2 (3.2)	1 (3)	1 (3.3)	
Severe	5 (7.9)	1 (3)	4 (13.3)	
Very severe	22 (34.9)	15 (45.5)	7 (23.3)	
DASS-Depression analysis, n (%)				
Normal	31 (49.2)	13 (39.4)	18 (60)	0.04
Mild	6 (9.5)	3 (9.1)	3 (10)	
Moderate	10 (15.9)	4 (12.1)	6 (20)	
Severe	6 (9.5)	6 (18.2)	0	
Very severe	10 (15.9)	7 (21.2)	3 (10)	
Scores of DASS-Stress, median (min-max)	6(0-18)	8 (0-18)	5(0-18)	0.01
Scores of DASS-Anxiety, median (min-max)	5(0-20)	7(0-20)	4(0-18)	0.07
Scores of DASS-Depression, median (min-max)	5(0-19)	7(0-18)	2(0-19)	0.01
Total scores of DASS-21, median (min-max)	18(0-54)	23(0-53)	10.5(0-54)	0.02
Fear of earthquake Scale, median (min-max)	25 (7-35)	27(7-35)	20.5(7-35)	0.04

DASS, Depression-Anxiety-Stress-Scale; IQR, interquartile range

Table III: Clinical characteristics of patients according to AE attacks status after earthquake

	Patients with worsened HAE during earthquake (n = 45)	Patients without (n = 18)	P value
Gender, n (%)			
Female	27 (60)	6 (33.3)	0.06
Male	18 (40)	12 (66.7)	
Age (years), median(min-max)	29 (18-77)	35 (21-63)	0.17
Body Mass Index, mean ± SD	26.6 ± 4.1	25.9 ± 4.04	0.53
HAE attack number, mean ± SD			
Pre-earthquake	4.7 ± 2.3	6.3 ± 3	0.02
Post-earthquake	10.2 ± 5.4	6.6 ± 3.6	0.01
HAE attack severity, median (min-max)			
Pre-earthquake	5 (1-10)	6.5 (2-8)	0.14
Post-earthquake	8 (5-10)	6 (2-8)	<0.001
Long term prophylaxis, n (%)			
Danazol	3 (6.7)	4 (22.2)	0.95*
Tranexamic acid	1 (2.2)	2 (11.1)	0.19*
C1 esterase inhibitor	0	0	0
Damage level in buildings, n (%)			
Intact	14 (32.6)	8 (44.4)	0.38
Slightly damage	21 (48.8)	10 (55.6)	0.63
Moderately damaged	5 (11.6)	0 (0)	0.31*
Severely damaged	3 (7)	0 (0)	0.55*
Collapsed	0 (0)	0 (0)	0
Death in the first degree relatives, n (%)	3 (6.7)	0 (0)	0.55*
Death in their acquaintances, n (%)	16 (35.6)	1 (5.6)	0.03*
Scores of DASS-Stress, median (min-max)	8 (0-18)	1.5 (0-9)	<0.001
Scores of DASS-Anxiety, median (min-max)	9 (0-20)	2 (0-12)	<0.001
Scores of DASS-Depression, median (min-max)	8 (0-19)	1 (0-10)	<0.001
Total scores of DASS-21, median (min-max)	23 (0-54)	6 (0-25)	<0.001
Fear of earthquake Scale, median (min-max)	28 (8-35)	12 (7-30)	<0.001

HAE, hereditary angioedema; IQR, interquartile range; SD, standard deviations; DASS, Depression-Anxiety-Stress-Scale

Table IV:The correlation analyses between psychological features and frequency and severity of HAE attacks

	Number of attacks		Severity of attacks	
	r	p	r	p
DASS-21 stress	0.517	<0.001	0.646	<0.001
DASS-21 anxiety	0.409	0.001	0.529	<0.001
DASS-21 depression	0.518	<0.001	0.624	<0.001
DASS-21 total	0.497	<0.001	0.628	<0.001
FES	0.488	<0.001	0.637	<0.001

DASS-21, Depression, Anxiety and Stress Scales-21; FES, Fear-of-Earthquake-Scale

Health conditions and medications after the earthquake

Three (4.8%) patients lost their first-degree relatives and 17 (27%) patients lost acquaintances in the earthquake.

The damage to the houses of the patients due to earthquakes was intact in 22 (34.9%), slightly damaged in 31 (49.2%), moderately damaged in 5 (7.9%), severely damaged in 3 (4.8%), and collapsed in 2 (3.2%). In the correlation analysis, the damage level in the buildings significantly correlated with the number of attacks and attack severity in the post-earthquake period (p:0.025, r:0.281 and p:0.042, r:0.257; respectively) (Figure 2).




Figure 2.The number and severity of HAE attacks after the earthquake according to the damage level in the buildings and FES score. Abbreviations: VAS, visual analogue scale; HAE, hereditary angioedema; FES, Fear-of-Earthquake-Scale

After the earthquake, 7 (11.1%) patients had problems in accessing icatibant. While 8 (12.7%) patients applied to the emergency department (ED), 14 (22.2%) patients did not go to the ED due to fear of earthquakes even though they needed it.

DISCUSSION

Until now, there have been no published studies assessing the impact of earthquake-induced mental health issues on the progression of HAE, making our findings significant in this regard. First, for about 70% and 50% of patients with HAE, the experience of the earthquakes exacerbated their physical and psychological symptoms, respectively. Second, about 10% of the patients with HAE experienced an interruption in medical welfare services following the earthquakes.

In the present study, patients with HAE experienced a significant increase in the number and severity of attacks after the earthquakes. While individuals with HAE often characterize most of their attacks as occurring without warning, stress emerges as the most commonly recognized triggering factor¹. In other research, stress emerged as the most commonly cited trigger, with trauma and infection following closely behind^{9,10}.

In our post-earthquake study, we observed a significant correlation between the number and severity of HAE attacks and psychological factors, including depression, anxiety, stress, and earthquake-related fear. These psychological parameters were assessed using two validated tools, namely DASS-21 and FES. In a study conducted by Karabacak et al., similar to our research, they reported positive correlations between the frequency and severity of HAE attacks and DASS-21 anxiety, stress, and depression scores, as well as the DASS-21 total scores, during the COVID-19 pandemic².

In our study, anxiety was observed in 58.7% of hereditary angioedema patients following the earthquake, while depression was detected in 50.8%. In patients with HAE, clinically significant anxiety symptoms have been reported by up to 15–46%, and relevant depressive symptoms have been observed in 18–42%^{11,12}. In the general population, the prevalence of depression after the Kumamoto earthquakes has been reported as 20.2%¹³. After conducting a systematic review and meta-analysis of studies pertaining on the 2010 Haiti earthquake, it was determined that the median prevalence of anxiety and depression in participants was 20.5% and 32.2%, respectively¹⁴. In our study, we were unable to examine the impact of pre-earthquake anxiety and depression statuses on the course of attacks as these were unknown for the patients. However, in our study, when compared with the literature, we observed elevated rates of anxiety and depression in measurements taken after the earthquake, both in comparison to typical HAE patients and to the post-earthquake population. In our study, the high prevalence of anxiety and depression among hereditary angioedema patients may be attributed to the fact that they already suffer from a life-threatening condition, which is further compounded by the occurrence of a natural disaster.

In our study, female patients exhibited significantly higher rates of anxiety, depression, and fear compared to their male counterparts following the earthquake. In the study by Christiansen et al., significantly higher rates of anxiety and depression were reported in women with HAE compared to men during the COVID-19 pandemic¹⁵. It is well-established that the prevalence of depression is higher in women than in men, even in non-disaster or non-pandemic contexts¹⁶. Furthermore, in the case of HAE, women have consistently exhibited a higher propensity for experiencing more

severe manifestations compared to their male counterparts, as reported in various studies¹⁷. However, in our study, while women exhibited a more pronounced deterioration in their HAE compared to men, this discrepancy did not achieve statistical significance.

In the research about the Great East Japan Earthquake, severe damage to the home and disruption of psychiatric care were reported as predictors of depressive symptoms after a big earthquake¹⁸. In our study, we observed a significant correlation between the level of damage in buildings and the number of attacks as well as their severity during the post-earthquake period. In addition to other earthquake victims, psychological and healthcare support should be provided particularly to HAE patients with severely damaged buildings.

The number of patients who lost access to their medications was lower than our initial expectations. It would serve patients with HAE well to always have at least a month of medicine available. The prompt regulatory measures implemented by the Ministry of Health to facilitate access to registered medications without requiring a prescription may have played an effective role. It is essential to keep in mind the importance of establishing consistent protocols for all earthquakes.

There are several limitations to this study. Firstly, the inability to assess the mental health status of our patients using identical scales during the pre-earthquake period limited our ability to draw more precise conclusions. Nevertheless, we believe that by excluding individuals who were on psychiatric medications or had psychiatric diagnoses, we have bolstered the robustness of our findings. Second, we could not compare our findings with a healthy control group exposed to the earthquake.

CONCLUSION

We identified the impacts of significant earthquakes on patients with HAE. The mental well-being of the patients was more susceptible to deterioration than their physical health.

In our study, we discovered that psychological factors, including anxiety, depression, stress, and fear, have the potential to exert a detrimental impact on the disease progression in individuals with HAE. This could potentially account for the heightened frequency of HAE attacks. As a result, it is recommended that individuals with HAE should be provided with early psychological support to address anxiety, stress, depression, and fear in the event of an earthquake.

Ethics Committee Approval: In adherence to the Declaration of Helsinki, the study received approval from the ethics committee at University of Health Sciences, Mehmet Akifinan Training and Research Hospital(Approval No:HRÜ/23.05.12, Date:27.03.2023). Informed consent forms were obtained from the patients.

Conflict of Interest: The authors declared no conflicts of interest.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Zotter Z, Csuka D, Szabo E, et al. The influence of trigger factors on hereditary angioedema due to C1-inhibitor deficiency. *Orphanet J Rare Dis.* 2014; 9:44.
2. Eyice Karabacak D, Demir S, Yeğit OO, et al. Impact of anxiety, stress and depression related to COVID-19 pandemic on the course of hereditary angioedema with C1-inhibitor deficiency. *Allergy.* 2021;76(8):2535-43.
3. Maurer M, Magerl M, Ansotegui I, et al. The international WAO/EAACI guideline for the management of hereditary angioedema-the 2017 revision and update. *Allergy.* 2018;73(8):1575-96.
4. Huskisson EC. Measurement of pain. *Lancet.* 1974;2(7889):1127-31.
5. Brown TA, Chorpita BF, Korotitsch W, Barlow DH. Psychometric properties of the Depression Anxiety Stress Scales (DASS) in clinical samples. *Behav Res Ther.* 1997;35(1):79-89.
6. Yılmaz Ö, Boz H, Arslan A. Depresyon Anksiyete Stres Ölçeğinin (DASS 21) Türkçe kısa formunun geçerlilik ve güvenilirlik çalışması. *Finans Ekonomi ve Sosyal Araştırmalar Dergisi.* 2017;2(2):78-91.
7. Akın A, Çetin B. Depresyon, Anksiyete Stres Ölçeği (DASÖ): Geçerlik ve güvenilirlik çalışması. *Kuram ve Uygulamada Eğitim Bilimleri.* 2007;7(1): 241-68.
8. Prizmić-Larsen Z, Vujčić MT, Lipovčan LK. Fear of COVID-19 and Fear of Earthquake: Multiple Distressing Events and Well-Being in Croatia. *Psychol Rep.*2023;332941231156813.
9. Alonso MLO, Valle SOR, Tortora RP, et al. Hereditary angioedema: a prospective study of a Brazilian single-center cohort. *Int J Dermatol.* 2020;59(3):341-4.
10. Jung JW, Suh DI, Park HJ, et al. Clinical features of hereditary angioedema in Korean patients: A nationwide multicenter study. *Int Arch Allergy Immunol.*2018;176(3-4):272-9.
11. Fouche AS, Saunders EF, Craig T. Depression and anxiety in patients with hereditary angioedema. *Ann Allergy Asthma Immunol.* 2014;112(4):371-5.
12. Caballero T, Aygören-Pürsün E, Bygum A, et al. The humanistic burden of hereditary angioedema results from the burden of illness study in Europe. *Allergy Asthma Proc.* 2014;35(1):47-53.
13. Kobayashi N. Psychosomatic medicine and disaster: experiences from the Kumamoto earthquake. *Jpn J Psychosom Med.* 2017;57:257-63.
14. Cénat JM, McIntee SE, Blais-Rochette C. Symptoms of posttraumatic stress disorder, depression, anxiety and other mental health problems following the 2010 earthquake in Haiti: A systematic review and meta-analysis. *J Affect Disord.*2020;273:55-85.
15. Christiansen SC, Lopes Veronez C, Smith TD, Riedl MA, Zuraw BL. Hereditary Angioedema: Impact of COVID-19 pandemic stress upon disease related morbidity and well-being. *Allergy Asthma Proc.*2023;44(2):115-21.
16. Albert PR. Why is depression more prevalent in women? *J Psychiatry Neurosci.* 2015; 40:219-21.
17. Busse PJ, Christiansen SC. Hereditary angioedema. *N Engl J Med.* 2020; 382:1136-48.
18. Tsuboya T, Aida J, Hikichi H, et al. Predictors of depressive symptoms following the Great East Japan earthquake: a prospective study. *Soc Sci Med.* 2016; 161:47-54.